

WILDBREW™ PHILLY SOUR

TECHNICAL DATA SHEET - WILDBREW™ SERIES

WILDBREW™ PHILLY SOUR

WildBrew™ Philly Sour is a unique species of *Lachancea* selected from nature by University of the Sciences in Philadelphia, PA, USA (Patent pending N° PCT/US20 18/043 148). WildBrew™ Philly Sour produces moderate amounts of lactic acid in addition to ethanol in one simple fermentation step. This first yeast in the WildBrew™ series is a great choice for innovative, sessionable sour beers with refreshing acidity and notes of stone fruit. With high attenuation, high flocculation and good head retention, WildBrew™ Philly Sour is an ideal yeast for traditional styles such as Berliner Weisse, Gose, American Lambic Style and American Wild Ales, and its resistance to hops make it perfect for Sour IPA's.



MICROBIOLOGICAL PROPERTIES

WildBrew™ Philly Sour is a pure strain of active dried yeast classified as *Lachancea* spp.

Typical Analysis of Philly Sour Yeast:

Percent solids	93% - 96%
Viability	≥ 1 x 10 ⁹ CFU per gram of dry yeast
Wild Yeast	WildBrew™ Philly Sour will grow on wild yeast media including Lysine, LCSM and LWYM
STA1 gene	Pure Strain is STA1 negative Contamination is undetectable by PCR test
Bacteria	<1 per 1 x 10 ⁶ yeast cells

Finished product is released to the market only after passing a rigorous series of tests

*See specifications sheet for details



BREWING PROPERTIES

In Lallemant's Standard Conditions Wort at 20°C (68°F) WildBrew™ Philly Sour yeast exhibits:

Fermentation that can be completed in 10 days.

High attenuation and High flocculation.

Aroma and flavor is sour, red apple and stone fruit, notably peach.


The optimal temperature range for WildBrew™ Philly Sour yeast when producing traditional styles is 20°C(68°F) to 25°C(77°F).

In the Lallemant standard conditions wort, typical pH of 3.2-3.5 and titratable acidity of 0.1-0.4% lactic acid are achieved. Higher lactic acid levels will be achieved in worts that are higher in glucose.

Lag phase, total fermentation time, attenuation and flavor are dependent on pitch rate, yeast handling, fermentation temperature and nutritional quality of the wort. *If you have questions please do not hesitate to contact us at brewing@lallemand.com*



FLAVOR & AROMA



QUICK FACTS

BEER STYLES

Berliner Weisse, Gose, lambic-style, American Wild, and Sour IPA

AROMA

Sour, stone fruit, peach and red apple

ATTENUATION

High

FERMENTATION RANGE

20°C(68°F) to 25°C(77°F)

FLOCCULATION

High

ALCOHOL TOLERANCE

9% ABV

PITCHING RATE

50 - 100g/hL to achieve a minimum of 0.5 - 1 million viable cells/mL



USAGE

The pitch rate will affect the fermentation performance and flavor of the beer. For WildBrew™ Philly Sour yeast, a pitch rate of 50-100 g per hL of wort is sufficient to achieve a minimum of 0.5 - 1 million viable cells/mL. More stressful fermentations such as high gravity, high adjunct or high acidity may require higher pitch rates and additional nutrients to ensure a healthy fermentation.

Standard pitch rate calculators may result in overpitching. Visit our Pitch Rate Calculator optimized for dry yeast samples at www.lallemandbrewing.com

WildBrew™ Philly Sour is a slow fermenting, killer negative strain and will tend to be outcompeted by other brewing strains. For this reason, it is not recommended to re-pitch this strain.

WildBrew™ Philly Sour is not recommended for bottle conditioning applications. A dedicated bottle conditioning strain such as LalBrew® CBC-1 should be used when bottle conditioning WildBrew™ Philly Sour fermentations.

As a non-*Saccharomyces* species, WildBrew™ Philly Sour should be treated according to normal best practices for handling wild yeast.



STORAGE

WildBrew™ Philly Sour yeast should be stored in a vacuum sealed package in dry conditions below 4°C (39°F). WildBrew™ Philly Sour will rapidly lose activity after exposure to air.

Do not use packs that have lost vacuum. Opened packs must be re-sealed, stored in dry conditions below 4°C (39°F), and used within 3 days. If the opened package is re-sealed under vacuum immediately after opening, yeast can be stored below 4°C (39°F) until the indicated expiry date. Do not use yeast after expiry date printed on the pack.

Performance is guaranteed when stored correctly and before the expiry date. However, Lallemand dry brewing yeast is very robust and some strains can tolerate brief periods under sub-optimal conditions.

If you have questions, do not hesitate to contact us. We have a team of technical representatives happy to help and guide you in your fermentation journey.



REHYDRATION

Rehydration of WildBrew™ Philly Sour in sterile water is recommended prior to pitching into wort in order to reduce stress on the cell as it transitions from dry to liquid form. For many fermentations, this stress is not significant enough to affect fermentation performance and flavor, so good results may also be achieved when pitching dry yeast directly into wort. We highly recommend rehydration in harsher fermentation conditions such as high gravity or sour wort where the added stress of dry-pitching is more likely to have a greater impact on the finished beer. Use of a rehydration nutrient such as Go-Ferm Protect Evolution has been shown to improve fermentation performance for difficult fermentations.

Rehydration guidelines are quite simple and present a much lower risk of contamination than a starter, which is unnecessary when using the recommended pitch rate of dried active yeast.

Sprinkle the yeast on the surface of 10 times its weight in clean, sterilized water at 30-35°C (86-95°F). Do not use wort, or distilled or reverse osmosis water, as loss in viability may result. **Stir gently**, leave undisturbed for 15 minutes, then stir to suspend yeast completely. Leave it to rest for 5 more minutes at 30-35°C.

Without delay, adjust the temperature to that of the wort by mixing aliquots of wort with the rehydrated yeast. Wort should be added in 5 minute intervals and taking care not to lower the temperature by more than 10°C at a time. Temperature shock of >10°C will cause formation of petite mutants leading to extended or incomplete fermentation and possible formation of undesirable flavors. Do not allow attemperation to be carried out by natural heat loss. This will take too long and could result in loss of viability or vitality.

Inoculate without delay into cooled wort in the fermenter. WildBrew™ Philly Sour yeast has been conditioned to survive rehydration. The yeast contains an adequate reserve of carbohydrates and unsaturated fatty acids to achieve active growth. It is unnecessary to aerate wort upon first use.

CONTACT US

For more information, please visit us online at www.lallemandbrewing.com

For any questions, you can also reach us via email at brewing@lallemand.com